

SUBSTITUTE FORM PTO-1350

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER
07258-022001

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371**

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)

09/700349

INTERNATIONAL APPLICATION NO.
PCT/EP99/03291INTERNATIONAL FILING DATE
May 12, 1999PRIORITY DATE CLAIMED
May 13, 1998

TITLE OF INVENTION

METHOD FOR PRODUCING PLANTS HAVING AN INCREASED TOLERANCE AGAINST DROUGHT AND/OR FUNGAL ATTACK AND/OR INCREASED SALT CONCENTRATIONS AND/OR EXTREME TEMPERATURE BY THE EXPRESSION OF PLASMODESMATA-LOCALIZED PROTEINS

APPLICANT(S) FOR DO/EO/US

ROHDE, Wolfgang, PRUFER, Dirk; TACKE, Eckhard; PASEMANN, Peter; SALAMINI, Francesco

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to promptly begin national examination procedures (35 U.S.C. 371(f)).
4. ☐ The US has been elected by the expiration of 19 months from the priority date (PCT Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☒ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☒ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11 to 16 below concern other documents or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☐ Other items or information:

- ☒ express mail declaration
☐ return postcard
☐
☐


CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No. EL584941327US

I hereby certify under 37 CFR §1.10 that this correspondence is being deposited with the United States Postal Service as Express Mail Post Office to Addressee with sufficient postage on the date indicated below and is addressed to the Commissioner for Patents, Washington, D.C. 20531


13 November 2000

Date of Deposit



Derek W. Norwood

Typed Name of
Person Signing

U.S. APPLICATION NO. (if known) 09/770549		INTERNATIONAL APPLICATION NO. PCT/EP99/03291	ATTORNEY'S DOCKET NUMBER 07258-022001
17. <input checked="" type="checkbox"/> The following fees are submitted: Basic National Fee (37 CFR 1.492(a)(1)-(5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1000 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$860 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$710 International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$690 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) \$100 ENTER APPROPRIATE BASIC FEE AMOUNT =			CALCULATIONS PTO USE ONLY
Surcharge of \$130 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).			\$860.00
Claims	Number Filed	Number Extra	Rate
Total Claims	- 20 =		x \$18
Independent Claims	- 3 =		x \$80
MULTIPLE DEPENDENT CLAIMS(S) (if applicable)			+ \$270
TOTAL OF ABOVE CALCULATIONS =			\$0.00
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.			\$430.00
SUBTOTAL =			\$430.00
Processing fee of \$130 for furnishing the English Translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f))			\$0.00
TOTAL NATIONAL FEE =			\$430.00
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +			\$0.00
TOTAL FEES ENCLOSED =			\$430.00
			Amount to be refunded:
			Charged:
a. <input checked="" type="checkbox"/> A check in the amount of \$430.00 to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. 06-1050 in the amount of \$0.00 to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 06-1050. A duplicate copy of this sheet is enclosed.			
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b) must be filed and granted to restore the application to pending status.			
SEND ALL CORRESPONDENCE TO:			
John R. Wetherell, Jr., Ph.D. FISH & RICHARDSON P.C. 4350 La Jolla Village Drive, Suite 500 San Diego, CA 92122 (858) 678-5070 phone (858) 678-5099 facsimile		 SIGNATURE: _____ NAME: John R. Wetherell, Jr., Ph.D. REGISTRATION NUMBER: 31,678	

09/700349

529 Rec'd PCT/PTC 14 NOV 2000

APPLICATION
FOR
UNITED STATES LETTERS PATENT

TITLE: METHOD FOR PRODUCING PLANTS HAVING AN
INCREASED TOLERANCE AGAINST DROUGHT AND/OR
FUNGAL ATTACK AND/OR INCREASED SALT
CONCENTRATIONS AND/OR EXTREME TEMPERATURE
BY THE EXPRESSION OF PLASMODESMATA-
LOCALIZED PROTEINS

APPLICANT: WOLFGANG ROHDE, DIRK PRUFER, ECKHARD TACKE,
PETER PASEMANN, AND FRANCESCO SALAMINI

CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No. EL584941327US

I hereby certify under 37 CFR §1.10 that this correspondence is being deposited with the United States Postal Service as Express Mail Post Office to Addressee with sufficient postage on the date indicated below and is addressed to the Commissioner for Patents, Washington, D.C. 20231.

November 13, 2000

Date of Deposit

Signature

Derek W. Norwood

Typed or Printed Name of Person Signing Certificate

8/PRTS

09/700349
529 Rec'd PCT/PTC 14 NOV 2000

Our Ref.: B 2729 PCT

Method for producing plants having an increased tolerance against drought and/or fungal attack and/or increased salt concentrations and/or extreme temperature by the expression of plasmodesmata-localized proteins

The invention relates to the use of nucleic acids which code for a (poly)peptide with an intrinsic affinity to plasmodesmata, to the production of plants or parts thereof having an increased tolerance against drought and/or fungal infections and/or increased salt concentrations; and/or extreme temperatures (heat/cold), and to corresponding methods. A plant, a plant tissue or a plant cell is advantageously transfected with the nucleic acid. The nucleic acid preferably codes for a virus-encoded transport protein which, in a particularly preferred embodiment, is a derivative of the pr17 protein with a hydrophilic N-terminal extension.

A number of documents are cited in this specification, the disclosure content of which is herewith incorporated by reference and the technical teaching of which can be applied within the meaning of the present invention.

One aim of classic plant breeding is the creation of productive varieties showing an increased tolerance against environmental factors or being resistant to stress factors. Those stress factors can be both of a biotic (insects, viruses, fungi etc.) and an abiotic nature (extreme temperatures, salt, drought). Whereas wildplants have adapted themselves to the extreme living conditions at stress-dominated habitats, drought, heat or salinity of the soil restrict the possibility of cultivating crop plants in such areas. On the other hand, agriculture also suffers heavy loss through abiotic stress at other habitats, as was shown in the USA in the year of drought in 1983: Almost half of the entire maize crop and a third of the expected soybean yield were destroyed as a consequence of persistent drought. All the cited abiotic stress factors impair the intercellular water balance. However, plants can adapt themselves to stress situations to a certain extent (Bohner, (1995) Plant Cell 7: 1099-1011). Proteins, for example, as well as metabolites of the plant metabolism like sugar alcohols, proline or glycine betaine have been identified as osmoregulators

* Translator's note: typographical error. Should probably read: "1099-1110".

accumulating as a result of abiotic stress. Based on that various strategies have been developed to produce transgenic plants having an increased tolerance against such factors or increased stress resistance by genetchnological modifications (article by Holmberg und Bülow (1998) Trends Plant Science 3: 61-66). An example of plant proteins as anti-stress factors are the so-called LEA (late embryogenesis abundant)-proteins the increased expression of which correlates with physiological and environmental pressure and which perform a protective function for the plant in extreme stress situations (see for example Chandler (1994) Annu. Rev. Plant Physiol. Plant Mol. Biol. 45: 113-141). The genetchnological modification of rice with the help of the barley-LEA-gene HVA1 indeed resulted in an increased tolerance against drought and salt (Xu (1996) Plant Physiology 110: 249-257). Further tests relating the expression of a LEA-genes in a heterologous system could not support these results (Iturriaga (1992) Plant Mol. Biol. 20: 555-558).

These plant metabolites identified as anti-stress factors or osmoprotectors include glycine betaine the effectiveness of which has been proved for example by maize plants (Saneoka, (1995) Plant Physiol. 107: 631-638). The synthesis of the glycine betaine in plant-chloroplasts is supported by the betaine aldehyde dehydrogenase (BADH) (Rhodes, (1993) Annu. Rev. Plant Physiol. Plant Mol. Biol. 44: 357-384). Transgenic tobacco plants expressing a bacterial BADH (Holmstrom, (1994) Plant. J. 6: 749-758) or a vegetable BADH (Rathinasabapathi, (1994) Planta 193: 155-162) showed the expected resistance to betaine aldehyde by conversion to glycine-betaine in amounts which are measured in plants exposed to stress. Nothing, however, was reported on an increased stress tolerance of these plants.

Obviously, derivatives of sugar like sugar alcohols or fructans are also produced and accumulated to a higher degree as a stress response by the plant. The increase of the mannitol level in transgenic tobacco plants by expression of a bacterial mannitol-1-phosphate-dehydrogenase resulted in an increased salt tolerance (Tarczynski, (1993) Science 259: 508-510). Equally, transgenic tobacco plants with a higher fructan level showed an increased resistance to drought compared to control plants (Pilon-Smith, (1995) Plant Physiol. 107: 125-130).

All the genetchnological methods mentioned up to now have an increased synthesis of osmoprotectors in the plant in common, by which the normal growth of the plant can be impaired (see Rathinasabapathi, op. cit.). In other cases bacterial genes are expressed in the plant, which is not necessarily accompanied by an optimal expression and therefore not by a suboptimal stress mastering. Consequently, the technical underlying of the present invention was to provide a method for creating

plants with an increased stress resistance which still show an essentially normal growth. This stress resistance (or tolerance) should preferably refer to both biotic and abiotic stress factors. This technical problem has been solved by providing the embodiments characterised in the claims.

Thus, the invention relates to the use of a nucleic acid which codes for a (poly)peptide with an intrinsic affinity to plasmodesmata to the production of plants or parts thereof having an increased tolerance against drought and/or fungal infections and/or increased salt concentrations and/or extreme temperatures, i.e. cold and/or heat. According to the invention, a plant, a plant tissue or a plant cell is usually transfected with such a nucleic acid according to standard methods.

All higher plants are characterised by their capability to create photosynthesis of sugars and their derivatives which can serve osmoprotectors in stress situations, as already mentioned above. The intercellular concentration of sugar and sugar derivatives - above all in the leaves for the protection of the photosynthetically-active chloroplasts - was solved, according to the invention, in an unexpectedly simple way by the measure mentioned above. Of a particular advantage is furthermore, apart from the fact that these measures can easily be worked out in a technical way by a skilled person that, according to the invention, the stress resistance can be increased with a mechanism which is valid in many and possibly even in all plants. Surprisingly, the plant tolerance against abiotic as well as against biotic factors could be increased with the method of the invention.

The term "increased salt concentration" refers to salt concentration in the soil leading to an increased ionic concentration in the plant which in turn leads to reduced growth. The absolute salt concentrations in the soil which have to be regarded as increased are different for different plants. They can, however, be determined by a skilled person according to standard methods, like for example by means of the disclosure content of Greenway and Munns (1980) *Ann. Rev. Plant. Physiol.* **31**: 149-190.

For the expression of the nucleic acid encoding a (poly)peptide with an intrinsic affinity to plasmodesmata in plant cells it is connected with regulatory sequences ensuring the transcription in plant cells. Among these are above all promoters. Generally, each promoter which is active in plant cells can be used for the expression.

The promoter can be either of a nature so that the expression takes place in a constitutive way or only in a certain kind of tissue, at a certain time of the plant development or at a point in time determined by external influences.

Concerning the plant the promoter can either be homologous or heterologous. Useful promoters are for instance the promoter of the 35S RNA of the Cauliflower Mosaic Virus (CaMV) and the ubiquitin-promoter of maize for a constitutive expression. Furthermore, there is usually a transcription-termination-sequence which serves the correct termination of the transcription and which can as well serve the addition of a poly-A tail to the transcript which performs a function regarding the stabilisation of the transcripts. Such elements like e.g. the terminator of the octopine synthetase gene of agrobacteria are described in the literature (see Gielen, EMBO J. 8 (1989), 23-29) and are interchangeable at will. In a preferred embodiment the promoter is the 35S CaMV promoter.

Apart from the nucleic acid which encodes a (poly)peptide with an intrinsic affinity to plasmodesmata the plant which is used or produced in any of the methods of the invention can contain further recombinant DNA-molecules which can, for example, be used for plant protection or quality enhancement of the plant or the harvest product thereof. Examples of plant protection measures include: (i) tolerance against herbicides (DE-A-3701623; Stalker (1988) Science 242, 419), (ii) resistance to insects (Vaek (1987) Plant Cell 5, 159-169), (iii) resistance to viruses (Powell (1986) Science 232, 738-743) and (vi)** resistance to ozone (Van Camp (1994) BioTech. 12, 165-168). Examples of quality enhancement are: (i) enhancement of imperishability of fruits (Oeller (1991) Science 254, 437-439), (ii) enhancement of the starch production in potato tubers (Stark (1992) Science 242, 419), (iii) modification of the composition of starch- (Visser (1991) Mol. Gen. Genet. 225, 289-296) and lipid compositions (Voelker (1992) Science 257, 72-74) and (iv) production of polymers foreign to plant (Porier (1992) Science 256, 520-523).

The availability of appropriate transformation systems is a prerequisite for the production of transgenic plants is. At the moment several methods are available for the transformation. The method for the transformation of dicotyledonous plants used most frequently is the gene transfer mediated by agrobacteria. In this connection the natural ability of the soil bacterium to integrate genetic material into the plant genome is used. Further appropriate methods are, for example, protoplast transformation by polyethyleneglycol-induced DNA-absorption, electroporation, sonication or microinjection as well as the transformation of intact cells or tissues through micro- or macroinjection in tissues or embryos, tissue electroporation, incubation of dry embryos in a solution containing DNA, vacuum infiltration of seeds and biolistic gene transfer (for an overall view: see e.g. Potrykus, Physiol. Plant (1990), 269-273 and Christou (1996) Trends in Plant Science 1, 423-431).

* Translator's note: typographical error. Should probably read: "iv"

Whereas the transformation of dicotyledonous plants via Ti-plasmid vector systems with the help of *Agrobacterium tumefaciens* is well established, more recent works indicate that monocotyledonous plants are indeed available for the transformation through vectors based on the *Agrobacterium* (Chan, *Plant Mol. Biol.* 22 (1993), 491-506; Hiei, *Plant J.* 6 (1994), 271-282; Bytner, *Proc. Natl. Acad. Sci. USA* 84 (1987), 5345-5349; Raineri, *Bio/Technology* 8 (1990), 33-38; Gould, *Plant Physiol* 95 (1991), 426-434; Mooney, *Plant, Cell Tiss. & Org. Cult.* 25 (1991), 209-218; Li, *Plant Mol. Biol.* 20 (1992), 1037-1048). The methods for the introduction of foreign DNA by means of the biolistic method or by protoplast transformation are known (see e.g. Christou (1996) *Trends in Plant Science* 1, 423-431; Willmitzer, L., 1993 *Transgenic plants*. In: *Biotechnology, A Multi-Volume Comprehensive Treatise* (H.J. Rehm, G. Reed, A. Pühler, P. Stadler, eds.) Vol. 2, 627-659, VCH Weinheim-New York-Basel-Cambridge). In the past it was possible to establish three of the above-mentioned transformation systems for various cereals: the electroporation of tissue, the transformation of protoplasts and the DNA-transfer through particle bombardment into regenerable tissues and cells (for an overall view: Jähne, *Euphytica* 85 (1995), 35 - 44). The transformation of wheat is described several times in the literature (for an overall view: Maheshwari, *Critical Reviews in Plant Science* 14 (2) (1995), 149 - 178). The skilled person can make use of the markers for the selection of transformed plant cells, plant tissues and plants. The transgenic plant cells can be regenerated into whole plants by means of techniques known to the skilled person. The skilled person can also make use of e.g. molecular biological methods like PCR in order to identify these plants. On the other hand, the skilled person can, of course, e.g. after self-fertilisation or back-crossing against the parent, also place seeds of such plants on selective media and by means of the germinative capacity of these seeds or the survival of the plants at a later stage of the development (depending on the chosen promoter) he can find out, whether the plants are transgenic or not. Basically, transgenic plants can be plants of any plant species, i.e. they can be either monocotyledonous or dicotyledonous plants.

As, regardless of the transformation method, only few cells have the desired characteristics a selectable marker is integrated into the plant genome in addition to the target gene in a conventional way. At the moment, above all genes which transmit tolerance towards herbicides or antibiotics are used for the selection of transformed plant cells. Appropriate resistance genes are, for example, the bar-gene of *Streptomyces hygroscopicus* which mediates resistance towards the total herbicide Phosphinothricin (De Block (1987) *EMBO J.* 6, 2513-2518) or the nptII-gene from the transposon Tn5 of *Escherichia coli* which leads to resistance towards

the antibiotic kanamycin (Herrera-Estrella (1983) EMBO J. 2, 987-995). Further selection systems are e.g. expression of a mannose-6-phosphate-isomerase and positive selection on nutritive media containing mannose (WO 94/20627). A further method makes use of the ability of a deaminase of *Aspergillus terreus* to detoxicate the insecticide blasticidin S (Tamura (1995) Biosci. Biotechnol. Biochem. 59, 2336-2338). Of course, the skilled person knows that the selection marker does not necessarily need to be present in the vector containing the recombinant DNA-molecule, but it can also be co-transformed with the vector (Lyznik (1989) Plant Mol. Biol. 13, 151-161; Peng (1995) Plant Mol. Biol. 27, 91-104). This possibility is, for example, useful if no physical linkage of the marker gene and the information which is to be transformed is desired.

In a preferred embodiment of the use according to the invention a plant is additionally regenerated from the transfected plant cell. The regeneration of the plant can take place according to methods which are conventional and known by the skilled person.

In an especially preferred embodiment according to the use of the invention further plants or plant cells are produced additionally of the regenerated plant after the regeneration step.

In a further preferred embodiment according to the use of the invention the (poly)peptide is a virus-encoded transport protein.

The expression of plant viral proteins which take part in the transport of viral information from cell to cell (transport proteins) also influence the transport or metabolism of starch, sugars and sugar derivatives in such a way that they accumulate to values exceeding the normal values during the light period in the photosynthetically active leaves of the plant (see e.g. Lucas 1993) *Planta* **190**: 88-96; Olesinski (1995), *Planta* **197**: 118-126; Olesinski, (1996) *Plant Physiol.* **111**: 541-550; Herbers (1997), *Plant J.* **12**: 1045-1056; Almon (1997), *Plant Physiol.* **115**: 1599-1607). The tolerance increase in of plants towards the above-mentioned environmental factors through expression of such proteins with an intrinsic affinity to plasmodesmata, the communicating galleries of adjoining cells in plants according to this preferred embodiment of the invention could not be deduced easily from the prior of the art. The normal function of virus-encoded transport proteins (TP) is to ensure the transport of the genetic information of a virus from cell to cell and thus to enable the spread of a virus from the original infection place to the whole plant.

in a particularly preferred embodiment of the use according to the invention the virus-encoded transport protein is the potato leaf roll virus-(PLRV) transport protein pr17 or a derivative thereof.

As illustrated in the examples the TP of the potato leaf roll-virus (PLRV) as well as the cultivated plant potato were chosen as a model system. The PLRV-TP which describes a characteristic of a particularly preferred embodiment of the method of the invention is a 17 kDa large protein (pr17) which is responsible for the transfer of the genomic RNA from cell to cell. It is encoded by means of the open reading frame (ORF) ORF4; this gene lies in the ORF3, the gene for the viral capsid protein CP, however, in a different reading frame. The protein possesses an aminoterminal domain for the formation of homopolymers (Tacke (1993), Virology 197: 274-282) and a carboxyterminal domain for binding single-stranded nucleic acids (Tacke (1991), J. Gen. Virol. 72: 2035-2038). This protein which is present in *in planta* phosphorylated form (Tacke (1993), op. cit.; Sokolova (1997), FEBS Lett. 400: 201-205) is expressed seven times more strongly than the viral-coat-protein gene (Tacke (1990), J. Gen. Virol. 71: 2265-2272). Both in PLRV-infected and pr17-transgene potato plants the pr17 is predominantly located in the plasmodesmata between sieve element and companion cell of the phloem (Schmitz (1997), Virology 235: 311-322), to which the virus is limited during its reproduction in the plant. Expression of a mutated pr17-protein in transgenic potato plants leads to broad-spectrum resistance against the most important potato viruses (Tacke (1996), Nature Biotechnology 14: 1597-1601). At the same time, however, in the course of this invention it was observed that the expression of WT and mutated PLRV-TPs in transgenic potato- and tobacco plants leads to an increase in the intracellular concentrations of sugars (sucrose, fructose, glucose) and derivatives of sugar like starch.

In a further preferred embodiment of the use according to the invention the derivative is a pr17 protein with a hydrophilic N-terminal extension.

Particularly good results in the sense of the invention were found when a variant of the pr17-gene was used which carries a N-terminal extension (pr17-N). In an embodiment the polylinker (multiple cloning site; MCS) of the Bluescript-vector was fused in a translational way to the 5'-end of the pr17-WT-gene and the first two AUG-translation codons of pr17 in ACG-codons were mutated by means of selective mutagenesis and an AUG-translation start codon was inserted into the polylinker sequence (Figur 1). This modification results in the expression of a derivative (pr17-N) of the pr17-WT-proteins with a hydrophilic extension through the sequence MAELGSGSELHRGGGRSRTS (SEQ ID NO: 1) at the amino terminus (Tacke et al., 1996, op. cit.; Figure 2). In the green-house test such transgenic potato plants show

broad-spectrum resistance against potato viruses PLRV, PVY and PVX as well as an increased concentration of sugar and sugar derivatives.

For the expression in plants this gene was brought under transcriptional control of the 35S-promoter and -terminator of the cauliflower-mosaic-virus (CaMV) in the vector pRT103 (Töpfer (1987), Nucleic Acids Res. 15: 5890) and this transcription unit (Figure 1) was then integrated into the binary plant transformation vector pBIN19 (Bevan (1984), Nucleic Acids Res. 12: 8711-8721). This vector was transferred to the *Agrobacterium tumefaciens* stem LBA4404 (pAL4404) (Hoekema (1983), Nature 303: 179-180) which was used for the transformation of *Solanum tuberosum* Var. Linda. Four (L4, L6, L7 and L3; see also Tacke (1996), op. cit.) of the independent transgenic potato lines as well as the initial potato variety Linda were chosen for further tests concerning induced tolerance.

As already mentioned above the hydrophilic extension comprises the amino acid sequence MAELGSGSELHRG:GGRSRTS in a particularly preferred embodiment of the use according to the invention.

In another embodiment of the use according to the invention the plant, the plant tissue or the plant cells stem from the potato, from tobacco, from cereals or vegetables or are potatoes, tobacco plants, cereal plants or vegetable plants.

In a further preferred embodiment of the use according to the invention the increase in the tolerance of the plants against fungal infections is a tolerance against infections with *Phytophthora infestans*.

As a surprising result according to this preferred embodiment it was found that transgenic lines received by the method according to the invention also distinguish themselves by a statistically significant tolerance against *Phytophthora infestans*, the pathogen of late blight of potato.

The invention further relates to the production of plants or parts thereof with an increased tolerance against drought and/or fungal infections and/or increased salt concentrations and/or extreme temperature (heat, cold), wherein

- (a) a plant, a plant tissue or a plant cell is transfected with a nucleic acid coding for a (poly)peptide with an intrinsic affinity to plasmadesmata.

Additionally, in a preferred embodiment of the method according to the invention

- (b) a plant is regenerated from the transfected plant cell.

In a particularly preferred embodiment of the method according to the invention

(c) further plants or plant cells are produced from the plant gained in (b) subsequent to step (b).

In a further preferred embodiment of the method according to the invention, the polypeptide is a virus-encoded transport protein.

In a particularly preferred embodiment of the method according to the invention the virus-encoded transport protein is the potato leaf roll virus-(PLRV) transport protein pr17 or a derivative thereof.

In a further preferred embodiment of the method according to the invention the derivative is a pr17 protein with a hydrophilic N-terminal-extension.

As already mentioned above in a particularly preferred embodiment of the method according to the invention the hydrophilic extension comprises the amino acid sequence MAELGSGSELHRGGGRSRTS.

In another embodiment of the method according to the invention the plant, the plant tissue or the plant cells stem from potato, from tobacco, from cereals or vegetables or are potatoes, tobacco plants, cereal plants or vegetable plants.

In a further preferred embodiment of the method according to the invention the increase in tolerance of the plant against fungal infections is a tolerance against infections with *Phytophthora infestans*.

The Figures show:

Figure 1

Production of the plasmid p17N. By specific mutagenesis the two AUG-codons of the wildtype pr17-gene were mutated to ACG and a translation initiation codon was inserted into the polylinker sequence.

Figure 2

Nucleotide and amino acid sequence of the mutated pr17N-gene or -protein.

Figure 3

Result of the resistance test #1 (in the green-house) with 5 plants each of the initial variety Linda (L) as well as the pr17N-transgenic lines L4, L6, L7 and L8.

A. Overall view of the test. B. View of one plant each per line.

Figure 4

Result of the resistance test #2: (in the phytochamber) with 6 plants each of the initial type Linda (L) as well as of the pr17N-transgenic lines L4, L6, L7 and L8.

A. Partial view of the overall test. B. View of selected plants.

Figure 5

Bonitation of the infestation of leaf disks in the laboratory test with *P. infestans* race 1-11 (test #2) after 9, 10 and 13 days after the infection (dpi).

Figure 6

Cumulative depiction of two tests concerning the infection of potato leaf disks with *P. infestans* race 1-11.

Figure 7

Habitus of the non-transgenic potato variety Linda (L) and of the 4 transgenic lines L4, L6, L7 and L8 after 5 weeks at 100 mM NaCl. Individual plants of the partial test (A) were described for a better depiction in (E). Concerning the initial variety Linda the lower leaves died off.

Figure 8

Habitus of individual plants of the non-transgenic potato variety Linda (L) and of the 4 transgenic lines L4, L6, L7 and L8 after 5 weeks with 100 mM NaCl. Concerning variety Linda there can be still recognised residues of the died off lower leaves as well as modifications of the stalk.

The Examples explain the invention.

Example 1: Production of the plasmid p17N

A modification at the 5'-end of the pr17-gene (ORF4) was achieved by translational fusion of the multiple cloning site of the Bluescript-vector, insertion of an optimised translation initiation codon as well as mutation of the two pr17-WT AUG initiation codon to ACG (Figure 1). This modification results in the expression of a derivative (pr17-N) of the pr17-WT-protein with a hydrophilic extension through the sequence MAELGSGSELHRGGGRSRTS at the amino terminus (Tacke (1996), op. cit.; Figure 2). The production of the plasmid p17N is described in Schmitz (1996), Nucleic Acids Res. **24**: 257-263 (therein named p17/NIII).

Example 2: Introduction of the T-DNA in the recipient organism

After the transformation of *E. coli* S17-1 cells and mating with the *A. tumefaciens* strain LBA 4404 (pAL 4404) (Hoekema *et al.*, 1983) agrobacteria carrying the plasmid p17N were used for plant transformation. Leaves of *S. tuberosum* var. Linda of the sterile culture were cut at the basis and incubated for 10 min in liquid MS-medium with an agrobacteria culture grown over night. After two days of incubation on solid MS-medium the leaves were washed and placed on selection-, regeneration-medium. This consisted of MS-medium complemented with 0.02 mg/l naphthyl acetic acid, 0.02 mg/l gibberellin acid, 2 mg/l zeatinriboside, 500 mg/l Claforan and 100 mg/l kanamycin sulfate. Sprouts developed after 6-8 weeks and were transplanted on MS-medium with 250 mg/l claforan and 150 mg/l canamycinsulfate for the root formation.

Example 3: Characterisation of the transgenic lines obtained

The expression of the N-terminal modified PLRV 17K TP was detected in Western blot with the help of 17K-specific antisera as described in Tacke *et al.*, 1996 (op. cit.). For this purpose protein extracts were produced of leaf material, the proteins were separated on a 12.5% polyacrylamide gel, transferred to nitrocellulose membranes and incubated with a 17K-specific antibody. The evidence of bound antibodies was carried out according to reliable methods by incubation with sheep-anti-rabbit-IgG/peroxidase conjugate and the peroxidase substrate of the ECL-kits (Amersham).

Example 4: Resistance test with *Phytophthora infestans*

For the determination of the quantitative resistance infestation tests were carried out in the laboratory using the pathogen of late blight of potato, *Phytophthora infestans*. The 1-11 race of the pathogen was used as inoculum (for the preservation this race is cultivated on leaf material of the potato variety Désirée). For this purpose small potato leaves were inoculated with the pathogen in an irrigation box (Gieffers (1989), J. Phytopathology 126: 115-132) which enables a permanent water supply, and were placed at 10°C. The light supply was effected by 10 lamps of the type Osram, L16, W/25 Weiß Universal for 16 h daily with a medium photoperiodically active radiation of about $100 \mu\text{mol s}^{-1}\text{m}^{-2}$. The sporangia which were formed after 8-10 days were washed off from the small leaves with water. The share of living sporangia was determined by the staining method according to Behr (Behr (1955) Zentralblatt für Bakt. Etc. II 108: 23/24, 641-656). For infestation tests a suspension with a sporangia density of 10^4 ml^{-1} was used.

Following the test according to Hodgson (Hodgson (1961), American Potato Journal 38: 259-264) a new testing method was developed. By means of a punching machine which was developed for the test, leaf disks of a diameter of 20 mm were produced. The punching method works with low pressure to avoid crushing of the leaf edges and necrosis as well as bacterial decomposition.

The leaf disks are placed in irrigation boxes (Glaeffers et al., op. cit.) on filtering paper with the bottom side up. The permanent water film on the filtering paper provides the leaf disks with sufficient water. For inoculation a drop with a sporangia suspension (200 sporangia/20 µl) is pipetted onto the middle of the leaf disk.

The inoculated leaf disks are incubated at a permanent temperature of 10°C and under the above-mentioned light conditions. Under these conditions the infectious zoosporangia hatch. After about 6 days the first sporangia developed, the infestation bonitation is effected after 8 to 10 days.

The leaf disk infestation, recognisable by sporangia turf-forming and leaf tissue decomposition, is assessed in percentage.

An important prerequisite for the test is that the potato plants which are to be tested are cultivated under the same conditions and that the leaf disks are gained from small leaves of the same leaf storey and the same leaf position.

In this way the quantitative infestation of green-house- and open land material can be tested. The degree of infestation decides on the quantitative degree of resistance.

Example 5: Induced tolerance against drought as an example of increased tolerance against abiotic stress

5 or 6 plants each of the transgenic lines L4, L6, L7 and L8 as well as the initial variety Linda were kept under drought stress for 8 weeks in the 6-8 leaf-stage with the plant being irrigated after 3 and 6 weeks. In this connection the transgenic plants showed a clearly increased tolerance against water stress in two independent tests as can be seen in Table 1 and the Figures 3 and 4. Table 1: Analysis of the potato lines L4, L6, L7 and L8 as well as the initial variety Linda after 8 weeks of water stress.

Line / type # surviving plants / # of the examined plant

	Test 1**	test 2***
Linda	1*/5	1*/6
L4	4/5	6/6
L6	5/5	6/6
L7	5/5	6/6
L8	5/5	5/6

*The surviving plant develops a new sprout from the tuber; all the original parts of the plant are dead in contrast to the surviving plants of the transgenic lines L4, L6, L7 and L8.

** This test was carried out under controlled conditions in the green-house.

***This test was carried out under controlled conditions in a phytochamber.

Example 6: Induced tolerance against *Phytophthora infestans* as an example of increased tolerance against fungal infection

Two independent infestation tests with *P. infestans* were carried out with the 1-11 races on leaf disks of green-house material. In this connection 8 leaf disks of 8 plants each per transgenic lines and initial variety Linda were used for infection in the laboratory test. The inoculum contained 300 sporangia per 20 µl water; the incubation of leaf disks was effected at 10°C and bonitations were carried out after 9, 10 and 13 days after the infection (dpi). All the 4 transgenic lines showed a significant lower infestation of up to 10 dpi compared to *P. infestans*. After that the infestation at L7 and L8 increased rapidly whereas L6 and particularly L4 maintained their relatively low degree of infestation (Figure 5). On average of both tests there is a significant difference regarding the resistance conduct of the 4 tested transgenic Linda-lines compared to the non-transgenic initial type Linda (Figure 6).

Example 7: Induced tolerance against salt as a further example of increased tolerance against abiotic stress

In a series of tests of 4 plants each of the lines L4, L6, L7 and L8 as well as the non-transgenic initial variety Linda (L) the potato plants were kept in granulate for 5

027003443-034504

weeks under salt stress, wherein salt in the form of NaCl in concentrations of 10, 25, 50, 75, 100, 200, 400 and 1000 mM was contained in the irrigatin water. The highest concentration of 1000 mM lead to heavy damage among all plants after 2 weeks and these plants were dead after 3 weeks, whereas the transgenic lines, however, not variety Linda, survived a salt concentration of 400 mM despite heavy damage. As depicted in Figures 7 and 8 for the irrigation with 100 mM NaCl after 5 weeks, all transgenic as well as all non-transgenic plants had a similar phenotype (the somewhat smaller growth height of the transgenic lines as depicted in Figure 7B is due to the genetic modification not to the salt stress): Concerning the non-transgenic initial variety Linda the lower leaves died off and the stalk shows characteristic pathological modifications (strong stenosis, brown coloring), as is also depicted in Figure 8, whereas regarding all of the transgenic lines the stalks are as green as before (Figure 7, 8) and the leaves did not die (Figure 7A,B). At most, there are a few isolated cases of necrosis on the lower leaves (Figure 8). Despite the salt stress all plants develop blossoms and tuber formation can be observed. (Figure 8).

Claims

1. Use of a nucleic acid encoding a (poly)peptide with an intrinsic affinity to plasmodesmata, for the production of plants or parts thereof having an increased tolerance against drought and/or fungal infections and/or increased salt concentration and/or extreme temperature (heat, cold).
2. The use of claim 1, wherein, further, a plant is regenerated from the transfected plant cell.
3. The use of claim 2, wherein, following the regeneration, additionally further plants or plant cells are produced from the regenerated plant.
4. The use of any one of claims 1 to 3, wherein the (poly)peptide is a virus-encoded transport protein.
5. The use of claim 4, wherein the virus-encoded transport protein is the potato leaf roll virus-(PLRV) transport protein pr17 or a derivative thereof.
6. The use of claim 5, wherein the derivative is a pr17-protein with a hydrophilic N-terminal extension.
7. The use of claim 5, wherein the hydrophilic extension is the amino acid MAELSGSGSELHRGGGRSRTS.
8. The use of any one of claims 1 to 7, wherein the plant, the plant tissue or the plant cells are derived from the potato, from tobacco, from cereal or vegetables or are potatoes, tobacco plants, cereal plants or vegetable plants.
9. The use of any one of claims 1 to 8, wherein the increase in tolerance of plants against fungal infections is an increase in tolerance against infections with *Phytophthora infestans*.
10. Method for producing plants or parts thereof having an increased tolerance against drought and/or fungal infections and/or increased salt concentrations and/or extreme temperature (heat, cold), wherein

(a) a plant, a plant tissue or a plant cell is transfected with a nucleic acid which encodes a (poly)peptide with an intrinsic affinity to plasmodesmata.

11. The method of claim 10, wherein further
(b) a plant is regenerated from the transfected plant cell.
12. The method of claim 11, wherein, further,
following step (b)
(c) further plants or plant cells are produced from the plant gained in
(b).
13. The method of any one of claims 10 to 12, wherein the (poly)peptide is a virus-encoded transport protein.
14. The method of claim 13, wherein the virus-encoded transport protein is the potato leaf roll virus-(PLRV) transport protein p17 or a derivative thereof.
15. The method of claim 14, wherein the derivative is a p17-protein with a hydrophilic N-terminal extension.
16. The method of claim 15, wherein the hydrophilic extension is the amino acid MAELSGSGSELHRGGGRSRTS:
17. The method of any one of claims 10 to 16, wherein the plant, the plant tissue or the plant cells are derived from potato, from tobacco, from cereals or vegetables or are potatoes, tobacco plants, cereal plants or vegetable plants.
18. The method of any one of claims 10 to 17, wherein the increase in tolerance of plants against fungal infections is an increase in tolerance against infections with *Phytophthora infestans*.

Abstract

The invention relates to the use of nucleic acids encoding a (poly)peptide with an intrinsic affinity to plasmodesmata, to the production of plants or parts thereof having an increased tolerance against drought and/or fungal infections and/or increased salt concentrations and/or extreme temperature (heat, cold), and to corresponding methods. A plant, a plant tissue or a plant cell is advantageously transfected with said nucleic acid. Preferably the nucleic acid encodes a virus-encoded transport protein which, in a particularly preferred embodiment, is a derivative of the pr17 protein with a hydrophilic N-terminal extension.





Fig. 1

[illegible]

Fig. 2

3/8



4/8

A



B





Fig. 4

Phytophthora infestans infestation on Linda etc.
Laboratory test, R 1-11



2nd test, October 1997

Figure 5

Linköping
Göteborg
Göteborg




Figure 6

Average of 2 tests, Aug. and Oct. 1997, 10 dpi

Wicoxon test: pillars with unequal letters differ from each other significantly at $P < 0.05$

7/8




Fig. 7

8/8



Fig. 8

INCREASED SAT
CALIZED PROTEINS

MAR 16 2001

PATENT & TRADEMARK OFFICE

ended by an agreement ref

Priority NOT Claimed

Pending

Paul N. Kokulis	16773	Kendrew H. Colton	30368	Roger R. Wise	31204	Anthony L. Miele	34393
G. Lloyd Kinghart	17698	G. Paul Edgell	24238	Michael R. Dzwonczyk	36787	Robert J. Walters	40862
Kevin E. Joyce	20508	Lynn E. Eccleston	35861	Jack S. Barufka	37087	Brian J. Beatus	38825
George M. Sirilla	18221	Adam A. Jakopin	32995	Adrian R. Hess	41835	John Jobe	28429
Donald J. Bird	25323	Mark G. Paulson	30793	William P. Atkins	38821	Mark C. Pickering	36239
Dale S. Lazar	28872	Stephen C. Glazier	31361	Paul L. Sharer	36004	David H. Jaffer	32243
Glenn J. Perry	28458	Richard H. Zaitlen	27248	Robin L. Teskin	35030	John R. Wetherell	31678
						Robert M. Bedonko	43488

PAT-116 301

**FOR UTILITY/DESIGN
CIP/PCT NATIONAL/PLANT
ORIGINAL/SUBSTITUTE/SUPPLEMENTAL
DECLARATIONS**

**RULE 63 (27 C.F.R. 1.63)
DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

**PW
FORM**

As a below named inventor, I hereby declare that my residence, post office address and citizenship are as stated below next to my name, and I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the **INVENTION ENTITLED**

the specification of which **(CHECK applicable BOX(ES))**
☒ A. ☐ is attached hereto.
☒ B. ☐ was filed on _____ as U.S. Application No. _____
☒ C. ☐ was filed as PCT International Application No. PCT/____ on _____
 and (if applicable to U.S. or PCT application) was amended on _____
 I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose all information known to me to be material to patentability as defined in 37 C.F.R. 1.56. Except as noted below, I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 356(f) of any foreign application(s) or patent or inventor's certificate, or 356(a) of any PCT International Application which designated at least one other country than the United States, listed below, and have also identified below any foreign application for patent or inventor's certificate, or PCT International Application, filed by me or my assignee disclosing the subject matter claimed in this application and having a filing date (1) before that of the application on which priority is claimed, or (2) as a priority claim, before the filing date of this application.

PRIOR FOREIGN APPLICATION(S)

Number	Country	Day/MONTH/Year Filed	Date First Laid-open or Published	Date Patented or Granted	Priority NOT Claimed
--------	---------	----------------------	-----------------------------------	--------------------------	----------------------

If more prior foreign applications, X box at bottom and continue on attached page.

As noted below, I hereby claim domestic priority benefit under 35 U.S.C. 119(e) or PCT international applications listed above or below, end, if this is a continuation-in-part (C/P) application. Insofar as the subject matter disclosed and claimed in this application is in addition to that disclosed in such prior applications, I acknowledge the duty to disclose all information known to me to be material to patentability as defined in 37 C.F.R. 1.56 which became available between the filing date of each such prior application and the national or PCT International filing date of this application.

PRIOR U.S. PROVISIONAL NONPROVISIONAL AND/OR PCT APPLICATION(S)

Application No. (series code/serial no.)	Day/MONTH/Year Filed	Status	Priority NOT Claimed
--	----------------------	--------	----------------------

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and I further declare that all statements were made with the knowledge that willful false statements are and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application of any patent issued thereon.

I and I hereby appoint Pillsbury Winthrop LLP, Intellectual Property Group, 1100 New York Avenue, N.W., Ninth Floor, East Tower, Washington, D.C. 20005-9618, telephone number (202) 861-0000 (to whom all communications are to be directed), and if below named persons (or the same address) individually and collectively my attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith and with the resulting patent, and I hereby authorize them to delete names/numbers below of persons no longer with their firm and to list and rely on instructions from and communicate directly with the persons/designations/firms/organization where/which first vend/sent this case to them as I by whom/which I hereby declare that I have consented after full disclosure to be represented unless/until I instruct the above firm and/or a below attorney in writing to the contrary.

Paul N. Kekulis	15272	Kandrow H. Colton	30089	Reyer R. Wise	31204	Anthony L. Miele	36393
G. Lloyd Knight	17898	G. Paul Edgall	30638	Michael P. Dzwonczyk	36787	Robert J. Walters	40862
Kevin E. Joyce	20509	Lynn E. Eccleston	35691	W. Patrick Bengtsson	32450	Brian J. Beatus	38525
George M. Sirilla	17821	Timothy J. Kilne	32854	Jar k S. Banfuka	37087		
Donald J. Bird	25923	David A. Jakorin	32995	Adam R. Hess	41885		
Dale S. Lazar	28872	Mark G. Paulson	30798	William P. Atkins	39621		
Paul E. White, Jr.	32011	Stephen C. Glazier	31361	Pa il L. Sharer	39004		
Siann J. Perry	28858	Richard H. Zeltner	27245	Rein L. Tockin	35030		

(1) INVENTOR'S SIGNATURE:

First		Middle Initial	Last	Date:	30.01.2001
Wolfgang			Kande		
Residence	Buseck	Germany	DEF	Germany	
City		State/Foreign Country		Country of Citizenship	
Mailing Address	Untergrasse 29				
(Include Zip Code)	35411				

(2) INVENTOR'S SIGNATURE:

First		Middle Initial	Last	Date:	6.02.2001
Dirk			Pfister		
Residence	Köln	Germany	DEF	Germany	
City		State/Foreign Country		Country of Citizenship	
Mailing Address	Hackländerstr. 15				
(Include Zip Code)	50825				

"X" box ☐ FOR ADDITIONAL INVENTORS, and proceed on the attached page to list each additional inventor.

☐ See additional foreign priorities on attached page (incorporated herein by reference).

Atty. Dkt. No. P

(MH)

DECLARATION AND POWER OF ATTORNEY

(continued)

ADDITIONAL INVENTORS

(3) INVENTOR'S SIGNATURE

Date: 02.02.2001

First		Middle Initial		Family Name	
Eckhard		Tacke			
Residence	First	Middle Initial		Family Name	
Post Office Address	City	State/Foreign Country		Country of Citizenship	
(include Zip Code)					

(4) INVENTOR'S SIGNATURE

Date: 31.01.2001

First		Middle Initial		Family Name	
Pekr		Fasemann			
Residence	First	Middle Initial		Family Name	
Post Office Address	City	State/Foreign Country		Country of Citizenship	
(include Zip Code)					

(5) INVENTOR'S SIGNATURE

Date: 31.01.2001

First		Middle Initial		Family Name	
Francesco		Salahini			
Residence	First	Middle Initial		Family Name	
Post Office Address	City	State/Foreign Country		Country of Citizenship	
(include Zip Code)					

(6) INVENTOR'S SIGNATURE

Date:

First		Middle Initial		Family Name	
Residence	First	Middle Initial		Family Name	
Post Office Address	City	State/Foreign Country		Country of Citizenship	
(include Zip Code)					

(7) INVENTOR'S SIGNATURE

Date:

First		Middle Initial		Family Name	
Residence	First	Middle Initial		Family Name	
Post Office Address	City	State/Foreign Country		Country of Citizenship	
(include Zip Code)					

(8) INVENTOR'S SIGNATURE

Date:

First		Middle Initial		Family Name	
Residence	First	Middle Initial		Family Name	
Post Office Address	City	State/Foreign Country		Country of Citizenship	
(include Zip Code)					

(9) INVENTOR'S SIGNATURE

Date:

First		Middle Initial		Family Name	
Residence	First	Middle Initial		Family Name	
Post Office Address	City	State/Foreign Country		Country of Citizenship	
(include Zip Code)					

PAT-1162 11/97